


Linka Lake

Pope County

Summary

Linka Lake is a moderately deep, mesotrophic lake. Algae concentration results (chlorophyll a) show that the lake experiences minor algae blooms occasionally in the summer. There are no detectable trends in water clarity over the past 20 years. This means that the lake is stable, with no indications of declining water quality. Linka Lake has a good amount of historical water quality monitoring data, which makes a lake evaluation like this possible. Monitoring should continue to enable future water quality analyses.



Lake Vitals

MN Lake ID:	61-0037-00
Ecoregion:	North Central Hardwood Forest
Major Watershed:	Chippewa River
Surface area (acres):	174.9
Littoral area (acres):	72.3
% Littoral area:	41.3%
Max depth:	50(ft) 15.2(m)
Aquatic Invasive Species:	None

Water Quality Characteristics


Years monitored: 1985 - 2016

Parameters	Historical	2016
Total Phosphorus Mean (ug/L):	26.0	21.8
Total Phosphorus Min (ug/L):	12	13
Total Phosphorus Max (ug/L):	64	29
Number of Observations:	107	5
Chlorophyll-a Mean (ug/L):	4.9	4.5
Chlorophyll-a Min (ug/L):	0.5	1.78
Chlorophyll-a Max (ug/L):	20	9.79
Number of Observations:	90	5
Secchi Depth Mean (ft):	13.6	16.7
Secchi Depth Min (ft):	2	10.6
Secchi Depth Max (ft):	28	20.6
Number of Observations:	316	5

Trophic State Index

Trophic State: Mesotrophic (46)


The figure below shows the minimum and maximum values with the arrows and the mean with the black dot (site 201).



Ecoregion Comparisons

(Primary site only. Comparisons are based on interquartile range, 25th - 75th percentile, for ecoregion reference lakes)

Ecoregion:	North Central Hardwood Forest
Total Phosphorus:	Within Expected Range
Chlorophyll-a:	Below Expected Range, which is better than expected
Secchi Depth:	Above Expected Range, which is better than expected




Trend Analysis Report

In assessing water quality, agencies and other lake data users want to know if the amount of algae has been changing over time. Scientists test hypotheses using statistics, and the hypothesis used in a trend analysis is that no trend exists. In other words, we begin with the assumption that there is no trend. We collect data and use statistics to determine the probability of collecting our data if this hypothesis of no trend is indeed true. The output from a statistical test is called the probability value (or p-value for short) of collecting data given the hypothesis of no trend is true. The smaller this probability value, the more likely the null hypothesis of no trend can be rejected. The MPCA has set the acceptable p-value to be less than 10%. In other words, if $p < 0.10$ we reject the hypothesis of no trend and accept that a trend likely exists. Another way to think of this is to say that there is in reality an existing trend, there is a 90% chance we would have collected the data we collected and that a 10% chance that the trend is a random result of the data. For detecting trends, a minimum of 8-10 years of data with four or more readings per season are recommended by the MPCA. Where data does not cover at least eight years or where there are only few samples within a year, trends can be misidentified because there can be different wet years and dry years, water levels, weather, and etc., that affect the water quality naturally.

Linka Lake had enough data to perform a trend analysis for all three parameters (Table 1). The data was analyzed using the Mann Kendall Trend Analysis.

Table 1. Trend analysis for Linka Lake.

Lake Site	Parameter	Date Range	Trend
202	Transparency	1994-2016	No Trend
202	Total Phosphorus	1996-2016	No Trend
202	Chlorophyll-a	1996-2002, 2004-2011, 2013-2016	No Trend



Linka Lake shows no evidence of water quality trends for any of the parameters monitored over the past 20 years. Overall, these trend results show that the water quality in Linka Lake is stable, with no indication of decline. Monitoring should continue so that this trend can be tracked in future years.